

Airborne fungal spore content in the atmosphere of the city of La Plata, Argentina

Andrea C. Mallo · Daniela S. Nitiu ·
María C. Gardella Sambeth

Received: 16 December 2009 / Accepted: 22 May 2010 / Published online: 8 June 2010
© Springer Science+Business Media B.V. 2010

Abstract The aim of this paper is to present the first aeromycological study of the atmosphere of the city of La Plata. Air samples were taken using a Hirst-type volumetric spore trap sampler (Lanzoni, VPPS 2000) for a period of a year (July 2000–June 2001). Seventy-nine morphological types of spores belonging to the Phyla Myxomycota, Zygomycota, Oomycota, Ascomycota, Basidiomycota, as well as anamorphs of Higher Fungi were identified. A total of 171670.21 spores were recorded, with a daily mean of 540 spores/m³. The spores were present throughout the year the study was carried out. However, there was a wide daily fluctuation in the concentration values with a tendency toward an increase during the summer months. The fungal spores were classified in three categories: *abundant*, having five spore types; *moderate* having 12 types; and *low*, with 62 morphological types which represent 67.2, 24.2, and 8.6%, respectively. The most representative taxa were *Cladosporium*

cladosporioides, *Leptosphaeria*, *Cladosporium herbarum*, *Coprinus*, and *Agaricus*, the first two taxa having a high frequency during the year of study. This scientific research reveals a great diversity of morphological types in this outdoor environment, showing a strong dominance of the Imperfect Fungi, whose components have seasonal rates. By this study, the first aeromycological profile of an urban center of Argentina has been portrayed, and new information on the field of aeromycology in the country has been provided.

Keywords Fungal spores · Diversity · Concentration · Frequency · Urban area · Argentina

1 Introduction

Fungi are cosmopolitan organisms which develop in the most varied substrates and live in virtually all climates in the earth. There is an important qualitative and quantitative variability in the anemophilous mycota of the various environments as well as local seasonal variations, having an impact on human health and air quality (Gregory 1961). These aerosols in particular tend to be especially difficult to study since their production, maturation, release, and eventual concentration of spores in the atmosphere depend on multiple biotic and abiotic local factors,

A. C. Mallo (✉) · D. S. Nitiu · M. C. Gardella Sambeth
Cátedra de Palinología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Calle 64 N°3, 1900 La Plata, Argentina
e-mail: malloa2001@yahoo.com.ar

A. C. Mallo
CIC, PBA, La Plata, Argentina

D. S. Nitiu · M. C. Gardella Sambeth
CONICET, Buenos Aires, Argentina

and on those resulting from human activity (Beggs 2004).

In Argentina, there have been few studies on the atmosphere mycobiota. Early aeromycological research in outdoor environments were carried out in the cities of La Plata (Iovannitti and Tiraboschi 1985; Negrin et al. 2007), Córdoba (Quiroga de Pascual and Nobile 1985); Resistencia (Mangiaterra et al. 1993), and Corrientes (Esquivel et al. 2003), by means of the gravitational method. The goals of these studies were the isolation, cultivation, and identification of taxa which were present in the atmosphere; these studies were oriented toward medical purposes. More recent research using Hirst-type samplers was done in the Alto Valle del Río Negro and Neuquén to analyze the presence, concentration, and impact of *Alternaria* spores on human health (Vega et al. 2001). Additionally, the diversity and dynamics of pollen and spores were also analyzed in San Carlos de Bariloche (Bianchi and Olabuenaga 2006). During the 1998–2001 period, an aerobiological monitoring was performed in the city of La Plata making use of the same method. In a first stage of the study, the diversity and concentration of pollen in the atmosphere, its relationship with meteorological features, as well as the pollen types which affect allergic respiratory diseases, was analyzed (Nitiu and Romero 2001, 2002; Nitiu and Mallo 2002; Nitiu 2003, 2004, 2006, 2008).

A second stage began with the analysis of the mycobiota, and its goal was to characterize the fungal portion, given the importance of these particles in the aerobiological record. The aim of this study was to provide the first identification and characterization of atmospheric mycobiota in La Plata by analyzing the diversity, concentration, representation, and frequency of the spores in the air during a year (July 2000–June 2001).

2 Materials and methods

2.1 Description of the area and location of the sampler

The city of La Plata (34°55'S, 57°57'W) is located in the northeast of the Province of Buenos Aires, Argentina. The region lies in the humid and subhumid mesothermal zone (Burgos and Vidal 1951) with a mean annual precipitation of 1,165 mm.

Temperature ranges from 2°C in June (mid - winter time) to 38.2°C in January (mid - summer time). The spore trap was placed on the top floor of a building situated at 15 m above ground level, near the geographic center of the city. The area is completely urbanized with green spaces and abundant trees.

2.2 Sampling technique

Over a total period of sampling (1998–2001), data from July 1, 2000 to June 14, 2001 were analyzed. Daily spore concentrations were sampled using a 7-day Hirst-type volumetric spore trap sampler Lanzoni model 2000 (Bologna, Italy). The flow rate was adjusted to 10 l/min. Spores were collected on Melinex adhesive tape and cut into daily segments (48 mm strips representing 24 h of exposure). These segments were then mounted on microscope slides in glycerol jelly containing fuchsine (0.1%) and protected with a cover glass. The spores were observed and identified using an Olympus BH2 microscope at a magnification of 400× along two longitudinal traverses. In order to achieve a more accurate identification, a magnification of 1000× was used in some cases. Spore counts were then converted into spores per cubic meter of air sampled (spore/m³ day) (Domínguez Vilches et al. 1991). Due to technical failures on the sampler, there was an interruption on data since 20/2/2001 to 8/3/2001 and 22/5/2001 to 29/5/2001.

2.3 Identification criteria

To characterize the different fungal types, the taxonomic and morphological classification of Saccardo (1886) was used. The descriptive terminology used is taken from the specialized bibliography (Barnet and Hunter 1987; Grant Smith 1990; Lacey and West 2006; Swartz 1971; Stearn 1973; Käärik et al. 1983; Ainsworth and Bisby 2001). Also, in the case of Myxomycota, the criteria of different authors (Aira et al. 2005; Lacey and West 2006; Docampo Fernández 2008) were taken into account.

2.4 Analysis criteria

According to the representativeness of each spore type in the total record, the following categories were considered: *abundant*, for those spore types exceeding

5% of the total concentration level; *moderate*, ranging from 1 to 4.99%; and *low*, below 0.99%.

Relative frequency of daily samples was calculated according to the criteria defined by Infante et al. (1991) and adapted by Herrero et al. (1995). The categories used were *very frequent* (75.1–100% of sampled days), *frequent* (36.6–75%), and *unfrequent* (0.1–36.5%).

3 Results

A total of 79 fungal taxa, which belong to Phyla Oomycota (3.8%), Zygomycota (3.8%), Myxomycota (1.26%), Basidiomycota (17.7%), Ascomycota (30.4%), and anamorphs of Higher Fungi (44.3%) were identified. They were classified according to their diagnostic characters (Table 1). Myxomycota spores were taken as a single category since it is very difficult to recognize with this methodology the numerous *taxa* which belong to this Phylum. The sampling performed resulted in a daily mean concentration of 540 spores/m³ and a total of 171670.21 spores for the period.

Figure 1 shows the constant presence of spores throughout the year, with a wide fluctuation in the daily values. During the year 2000, there was a daily mean of 284.51 spores/m³ in winter (June–September), and a mean of 394.98 spores/m³ in spring (September–December), with maximum values that did not exceed 2,000 spores/m³ per day. Summer 2001 (December–March) showed an increase in the fluctuations and concentrations rose to a mean value of 814.43 spores/m³ per day, with a peak of 3478.53 spores/m³ on 17 February 2001. In autumn (March–June), there was a slight decrease in mean daily values with 714.96 spores/m³. However, the maximum annual concentration value was recorded on 7 June, with a result of 4753.61 spores/m³.

A study of the monthly distribution of spore concentration in their respective Phyla (Oomycota, Myxomycota, Zygomycota, Ascomycota, Basidiomycota, and Imperfect Fungi) was carried out; a different dynamics for each group was observed. For example, a combined analysis of the first three Phyla showed that the total maximum monthly concentration of all spore types (1175.93) was detected in December, with a greater occurrence of Myxomycota (Fig. 2). April showed a high presence of spores, being the month when Zygomycota and Oomycota reached

their peak. On the other hand, the minimum monthly concentration (110.49) was registered in October. Spores belonging to Ascomycota, Basidiomycota, and Imperfect Fungi (Fig. 2) were present in high concentrations throughout the year, the monthly total maximums were observed during the period going from December to March. In February, the highest spore concentration (20278.09) was recorded; 69.35% of which were asexual spores. As for diversity, the Imperfect Fungi contributed especially with *Cladosporium*, *Alternaria*, and *Periconia* spores throughout the year, their peak occurring in February. The Phylum Basidiomycota is mainly represented by *Coprinus*, *Agaricus*, and *Agrocybe*, and their maximum concentration values coincided with the previous group. Ascomycota spores mostly corresponded to *Leptosphaeria* and other related phragmospores, being June the month with the highest concentration.

The analysis of representativeness (Table 1) revealed the existence of five morphological types in the *abundant* category accounting for 67.2% of the total record. *Cladosporium cladosporioides*, *Leptosphaeria*, *Cladosporium herbarum*, *Coprinus*, and *Agaricus* were the dominant types. Among *moderates*, 12 *taxa* were identified, most of them conidia, contributing with 24.2% of the total spectrum. The most significant ones were *Alternaria*, *Aspergillus/Penicillium*, *Periconia*, and *Agrocybe*. In the *low* category, 62 spore types were registered which accounted for 8.6% of the total. There is a wide variety of morphologies and components of both sexual and asexual origin, being *Monodictis* and *Didymella* the most prominent.

Frequency analysis of fungal spores in the atmosphere (Table 1) revealed the presence of six *very frequent* types: *Leptosphaeria* (91.4%), *Cladosporium herbarum* (86.4%), *C. cladosporioides* (84.8%), *Alternaria* (81.6%), *Agrocybe* (80.7%), and *Agaricus* (80.4%). In the *frequent* category, there were 28 types; the most outstanding representatives were *Periconia* (74.4%), *Arthrinium* (71.2%), and *Coprinus* (70%). In the *unfrequent* category, 45 types were found, among which *Polythrincium* (34.8%) and *Monodictis* (34.2%) were dominant.

4 Discussion

Consistent with observations in other parts of the world, the atmosphere of the city of La Plata has a

Table 1 Fungal spore types identified in the atmosphere of La Plata during July 2000–June 2001

Phylum	Spore type	Morphological type	Percentage of representation	Category	Frequency (% days)
Oomycota	<i>Peronospora</i>	Hyalospore	1.0750	M	**
	<i>Plasmopara</i>	Hyalospore	0.0164	L	*
Zygomycota	<i>Absidia</i>	Hyalospore	0.5100	L	*
	<i>Cunninghamella</i>	Hyalospore	0.0900	L	*
	<i>Ryzyopus</i>	Hyalospore	0.0800	L	*
Ascomycota	<i>Amphisphaeria</i>	Phaeodidymospore	0.0712	L	*
	<i>Caloplaca</i>	Hyalospore	0.0164	L	*
	<i>Chaetomium</i>	Phaeospore	0.4081	L	**
	<i>Chaetosphaerella</i>	Phaeophragmospore	0.0053	L	**
	<i>Cryphonectria</i>	Hyalodidymospore	0.0800	L	**
	<i>Diaporthe</i>	Hyalodidymospore	0.0890	L	*
	<i>Diatrypaceae</i>	Hyaloscoleospore	0.0266	L	*
	<i>Didymella</i>	Hyalodidymospore	0.5100	L	**
	<i>Didymosphaeria</i>	Phaeodidymospore	0.4178	L	**
	<i>Leptosphaeria</i>	Hyalophragmospore	8.1953	A	***
	<i>Lophiostoma</i>	Phaeophragmospore	0.7965	L	*
	<i>Massariosphaeria</i>	Phaeophragmospore	0.4425	L	*
	<i>Melanomma</i>	Phaeophragmospore	0.8850	L	*
	<i>Mycosphaerella</i>	Hyalodidymospore	0.5310	L	**
	<i>Paraphaeosphaeria</i>	Phaeophragmospore	1.0050	M	**
	<i>Pleospora</i>	Phaeodictyospore	1.2730	M	*
	<i>Saccobolus</i>	Phaeodidymospore	0.0091	L	*
	<i>Sporormiella</i>	Phaeodidymospore	0.1386	L	*
	<i>Sordaria</i>	Phaeospore	0.2765	L	*
	<i>Trichocladium</i>	Phaeodidymospore	0.0996	L	*
	<i>Venturia</i>	Hyalodidymospore	0.0052	L	*
	<i>Xylariaceae</i>	Phaeospore	1.0252	M	**
	Anamorph	<i>Alternaria</i>	Phaeodictyospore	3.8372	M
<i>Arthrimum</i>		Phaeospore	1.7976	M	**
<i>Aspergillus/Penicillium</i>		Hyalospore	3.8821	M	**
<i>Bipolaris</i>		Phaeophragmospore	0.6315	L	**
<i>Botrytis</i>		Hyalospore	0.2152	L	*
<i>Briosia</i>		Hyalarthrospore	0.0267	L	*
<i>Cercospora</i>		Hyaloscoleospore	0.2334	L	*
<i>Cerebella</i>		Phaeodictyospore	0.0195	L	*
<i>Cladosporium cladosporioides</i>		Hyalospore	31.7595	A	***
<i>Cladosporium herbarum</i>		Hyalospore	11.4551	A	***
<i>Curvularia</i>		Phaeophragmospore	0.3181	L	**
<i>Epicoccum</i>		Phaeodictyospore	0.6547	L	**
<i>Exosporium</i>		Phaeophragmospore	0.1653	L	**
<i>Exserohilum</i>		Phaeophragmospore	0.0826	L	**
<i>Fusarium</i>		Hyalophragmospore	0.3114	L	*

Table 1 continued

Phylum	Spore type	Morphological type	Percentage of representation	Category	Frequency (% days)
	<i>Fusicladium</i>	Hyalodidymospore	0.0100	L	*
	<i>Geotrichum</i>	Hyalarthrospore	0.0267	L	*
	<i>Gliomastix</i>	Phaeospore	0.0014	L	*
	<i>Helicomycetes</i>	Sympodulospore	0.0125	L	*
	<i>Helminthosporium</i>	Phaeophragmospore	0.2479	L	**
	<i>Monodictis</i>	Phaeodictyospore	0.7942	L	*
	<i>Nigrospora</i>	Phaeospore	0.0699	L	*
	<i>Ochroconis</i>	Phaeodidymospore	0.1139	L	**
	<i>Oidiodendron</i>	Hyalarthrospore	0.0106	L	*
	<i>Oidium</i>	Hyalospore	1.6244	M	*
	<i>Paecilomyces</i>	Hyalospore	0.0190	L	*
	<i>Periconia</i>	Phaeospore	3.3005	M	**
	<i>Pestalotiopsis</i>	Phaeophragmospore	0.0012	L	*
	<i>Pithomyces</i>	Phaeodictyospore	0.2169	L	**
	<i>Podosporium</i>	Phaeophragmospore	0.0025	L	*
	<i>Polythrincium</i>	Hyalodidymospore	0.3055	L	**
	<i>Pseudocercospora</i>	Phaeoscolecospore	0.0496	L	*
	<i>Spegazzinia</i>	Phaeostaurospore	0.0509	L	*
	<i>Sporidesmium</i>	Phaeophragmospore	0.0826	L	**
	<i>Stemphylium</i>	Phaeodictyospore	0.2063	L	**
	<i>Tetraploa</i>	Phaeostaurospore	0.0202	L	*
	<i>Torula</i>	Phaeophragmospore	0.9960	M	**
Basidiomycota	<i>Agaricus</i>	Phaeospore	5.1937	A	***
	<i>Agrocybe</i>	Phaeospore	2.7537	M	***
	<i>Coprinus</i>	Phaeospore	6.8508	A	**
	<i>Cortinarius</i>	Phaeospore	0.3376	L	*
	<i>Ganoderma</i>	Hyalospore	0.6220	L	**
	<i>Gymnopilus</i>	Phaeospore	0.0924	L	*
	<i>Inocybe</i>	Phaeospore	0.0007	L	*
	<i>Lepiota</i>	Hyalospore	0.0430	L	*
	<i>Lycoperdon</i>	Phaeospore	0.0307	L	*
	<i>Puccinia</i>	Phaeospore	0.3132	L	*
	<i>Psathyrella</i>	Phaeospore	0.0228	L	*
	<i>Tilletia</i>	Phaeochlamydospore	0.1608	L	*
	<i>Uredinales</i>	Hyalospore	0.4102	L	**
	<i>Ustilaginales</i>	Phaeospore	0.0758	L	*
Myxomycota		Phaeospore	1.1853	M	**

Representativeness (A abundant; M moderate; L low). Frequency: (***) very frequent, ** frequent and * unfrequent)

wide diversity of spore types with a significant contribution of mitotic spores and elements belonging to Oomycota, Myxomycota, Zygomycota, Ascomycota, and Basidiomycota (Aira et al. 2006; Calderón

et al. 1997; De Antoni Zoppas et al. 2006; Díez Herrero et al. 2006; Fernández et al. 1998; Grinn-Gofrón and Rapiejko 2009; Ibañez Henríquez et al. 2001; Stepalska and Wolek 2005).

Fig. 1 Total daily concentration of fungal airborne spores in La Plata during July 2000–June 2001





Fig. 2 Monthly concentration of anemophilous spores belonging to the Phyla Oomycota, Zygomycota, Myxomycota, Ascomycota, Basidiomycota, and Anamorphic Types



This study revealed a greater diversity in comparison with studies carried out in other cities of Argentina such as: Resistencia, Corrientes, and

Córdoba (Mangiaterra et al. 1993; Esquivel et al. 2003; Quiroga de Pascual and Nobile 1985); similar results were obtained from prior aerobiological

studies made in the city of La Plata (Iovannitti and Tiraboschi 1985; Negrin et al. 2007). However, it is difficult to make comparisons in terms of diversity and spore concentrations due to the different methodologies used in the area; it is therefore advisable to use combined both, viable and no viable volumetric methods.

Quantitative seasonal variation shows an increase in concentrations as the year goes by, and peaks occur in summer and late fall. It was observed that in moderate climates the highest spore concentrations occur during this period (Kasprzyk 2008).

Asexual spores are dominant in the total record. Among them, *Alternaria* and *Cladosporium* are highly abundant throughout the year, reaching their peak, together with the Basidiomycota spores, during summer.

The most abundant spores are also characterized by their frequency during the year. Thus, *Leptosphaeria* and *Cladosporium* in their two forms (*C. cladosporioides* y *C. herbarum*) occur almost throughout the period. Previous spore types are common anemophilous components of the atmosphere of La Plata and other cities worldwide (Horner et al. 1995; Cruz et al. 1997; Myskowska et al. 2002; Corden et al. 2003).

5 Conclusions

Fungal spores are present throughout the year in the atmosphere but their diversity and concentration show seasonal fluctuations. Despite their great diversity, only five morphological types are the main contributors to the studied record, and these fungal spores contain allergens involved in respiratory diseases (Simmon Nobbe et al. 2008). Imperfect Fungi spores are prevalent in the warmer and dry months. On the contrary, in cool periods of high relative humidity, highest concentrations of Ascomycota are detected.

By this study, the first aeromycological profile of an urban center of Argentina has been portrayed, and new information on the field of aeromycology in the country has been provided.

Acknowledgments This study was carried out with the financial support of Universidad Nacional de La Plata, Consejo Nacional de Investigaciones Científicas y Tecnológicas

(CONICET) and Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CIC - PBA). We also thank the reviewers for their recommendations and suggestions.

References

- Ainsworth, G. C., & Bisby, G. R. (2001). *Dictionary of the fungi* (9th ed.). Kew: Commonwealth Mycol. Inst.
- Aira, M. J., Jato, V., & Iglesias, I. (2005). Calidad del aire. Ed. Xunta de Galicia.
- Aira, M. J., Rodríguez Rajo, F. J., & Jato, V. (2006). Comportamiento temporal de las mitosporas de *Cladosporium* en la atmósfera de Galicia (España). *Boletín Micológico*, 21, 19–25.
- Barnet, H. L., & Hunter, B. B. (1987). *Illustrated genera of imperfect fungi*. New York: MacMillan Publ. Co.
- Begg, P. J. (2004). Impacts of climate change on aeroallergens: Past and future. *Clinical and Experimental Allergy*, 34, 1507–1513.
- Bianchi, M. M., & Olabuenaga, S. E. (2006). A 3-year airborne pollen and fungal spores record in San Carlos de Bariloche, Patagonia, Argentina. *Aerobiologia*, 22, 247–257.
- Burgos, J. J., & Vidal, A. L. (1951). Los climas de la República Argentina según la nueva clasificación de Thornthwaite. *Meteoros*, 1(1), 3–32. Buenos Aires.
- Calderón, C., Lacey, J., Mc Cartney, A., & Rosas, I. (1997). Influence of urban climate upon distribution of airborne Deuteromycete spore concentration in Mexico City. *International Journal of Biometeorology*, 40, 71–80.
- Corden, J. M., Millington, W. M., & Mullins, J. (2003). Long-term trends and regional variation in the aeroallergen *Alternaria* in Cardiff and Derby, UK—are differences in climate and cereal production having effect? *Aerobiologia*, 19, 191–199.
- Cruz, A., Saenz de Santamaría, M., Martínez, J., Martínez, A., Guisantes, J., & Palacios, R. (1997). Fungal allergens from important allergenic fungi imperfecti. *Allergologia et Immunopathologia*, 25(3), 153–158.
- De Antoni Zoppas, B. C., Valencia Barrera, R. M., Vergamini Duso, S. M., & Fernández-González, D. (2006). Fungal spores prevalent in the aerosol of the city of Caxias do Sul, Rio Grande do Sul, Brazil, over a 2-year period (2001–2002). *Aerobiologia*, 22, 119–126.
- Díez Herrero, A., Sabariego Ruíz, S., Gutierrez Bustillo, M., & Cervigón Morales, P. (2006). Study of airborne fungal spores in Madrid, Spain. *Aerobiologia*, 22, 135–142.
- Docampo Fernández, S. (2008). Estudio Aerobiológico de la atmósfera de la costa oriental de Málaga (sur de España) e incidencia de las esporas fúngicas en el interior de la Cueva de Nerja. Tesis Doctoral. Facultad de Ciencias. Universidad de Málaga. España.
- Domínguez Vilches, E., Galán, C., Villamandos, F., & Infante, F. (1991). Handling and evaluation of the data from the aerobiological sampling. *Monografías REA/EAN*, 1, 1–18.
- Esquivel, P., Mangiaterra, M., Giusiano, G., & Sosa, M. (2003). Microhongos anemófilos en ambientes abiertos de dos ciudades del nordeste argentina. *Boletín Micológico*, 18, 21–28.

- Fernández, D., Valencia, R. M., Molnár, T., Vega, A., & Sagües, E. (1998). Daily and seasonal variations of *Alternaria* and *Cladosporium* airborne spores in León (North–West, Spain). *Aerobiologia*, *14*, 215–220.
- Grant Smith, E. (1990). *Sampling and identifying allegenic pollens and molds*. San Antonio, Texas: Blewstone Press.
- Gregory, P. H. (1961). *The microbiology of the atmosphere*. Leonard Hill Limited (London). New York: Interscience Publishers, Inc.
- Grinn-Gofrón, A., & Rapiejko, P. (2009). Occurrence of *Cladosporium* spp. and *Alternaria* spp. spores in Western, Northern and Central-Eastern Poland in 2004–2006 and relations to some meteorological factors. *Atmospheric Research*. doi:10.1016/2009.02.014.
- Horner, W. E., Helbling, A., Salvaggio, J. E., & Leher, S. B. (1995). Fungal allergens. *Clinical Microbiology Reviews*, *40*, 161–179.
- Ibañez Henriquez, V., Rojas Villegas, G., & Roure Nolla, J. M. (2001). Airborne fungi monitoring in Santiago, Chile. *Aerobiologia*, *17*, 137–142.
- Iovannitti, C., & Tiraboschi, I. N. (1985). Hongos anemófilos de la ciudad de Plata y preparación de sus antígenos. *Revista Argentina de Micología*, *8*(2), 6–12.
- Käärik, A., Keller, J. Kiffer, E., Perreau, J., & Reisinger, O. (1983). Atlas of airborne fungal spores in Europe. In S. Nilsson (Ed.). Berlin: Springer-Verlag.
- Kasprzyk, I. (2008). Aeromycology—Main research fields of interest during the last 25 years. *Annals of Agricultural and Environmental Medicine*, *15*, 1–7.
- Lacey, M. E., & West, J. S. (2006). *The air spora*. Dordrecht, The Netherlands: Springer.
- Mangiaterra, M., Alonso, J. M., Medina, E., & Cerbera, L. (1993). Micoflora Anemófila de la ciudad de Resistencia. *Revista Argentina de Micología*, *16*, 10–16.
- Myskowska, D., Stepalska, D., Obtulowicz, K., & Porebski, G. (2002). The relationship between airborne pollen and fungal spore concentration and seasonal pollen allergy symptoms in Cracow in 1997–1999. *Aerobiologia*, *18*, 153–161.
- Negrin, M. M., Del Panno, T., & Ronco, A. E. (2007). Study of bioaerosols and site influence in the La Plata area (Argentina) using conventional and DNA (fingerprint) based methods. *Aerobiologia*, *23*, 249–258.
- Nitiu, D. S. (2003). Annual, daily and diurnal variations of *Celtis* airborne pollen in La Plata (Argentina). *Aerobiologia*, *19*(2), 71–78.
- Nitiu, D. S. (2004). Intradiurnal fluctuation pollen in La Plata. Argentina. Part I. Herbaceous taxa. *Aerobiologia*, *20*, 69–74.
- Nitiu, D. S. (2006). Aeropalinologic analysis of La Plata city (Argentina) during a 3-years period. *Journal International of Aerobiology*. *Aerobiologia*, *22*, 79–87.
- Nitiu, D. S. (2008). Estudio aeropalinológico de la ciudad de La Plata. *Tesis Doctoral* 888. Biblioteca Florentino Ameghino Inv. T 1819. *Facultad de Ciencias Naturales y Museo*. UNLP.
- Nitiu, D. S., & Mallo, A. (2002). Incidence of allergenic pollen of *Acer* spp., *Fraxinus* spp. and *Platanus* spp. in the city of La Plata, Argentina: preliminary results. *Aerobiologia*, *18*, 65–71.
- Nitiu, D. S., & Romero, E. (2001). Contenido polínico de la atmósfera de la ciudad de la Plata, Argentina. *Polen*, *11*, 79–85.
- Nitiu, D. S., & Romero, E. (2002). Caracterización aeropalinológica de la atmósfera de la ciudad de la Plata. Vinculación con alergias respiratorias. *Boletín Sociedad Argentina Botanical*, *37*(1–2), 79–85.
- Quiroga de Pascual, R. L., & Nobile, R. (1985). Incidencia de hongos ambientales durante un año en la ciudad de Córdoba. *Revista Argentina de Micología*, *8*, 16–22.
- Saccardo, P. A. (1886). *Sylloge fungorum*, *4*, 1–8. Pavia.
- Simmon Nobbe, B., Denk, U., Pöll, V., Rid, R., & Breitenbach, M. (2008). The spectrum of fungal allergy. *International Archives of Allergy and Immunology*, *145*, 58–86.
- Stearn, W. T. (1973). *Botanical Latin* (2nd ed.). Newton Abbot: David and Charles.
- Stepalska, D., & Wolek, J. (2005). Variation in fungal spore concentrations of selected taxa associated to weather conditions in Cracow, Poland in 1997. *Aerobiologia*, *21*, 43–52.
- Swartz, D. (1971). *Collegiate dictionary of botany*. New York: Ronald Press Co.
- Vega, L., Bianchi, M. M., & Nordstrom, G. (2001). *Alternaria* and allergy in the Río Negro and Neuquén Upper Valley, North Patagonia, Argentina: A preliminary approach. *Allergy*, *54*, 15.