

Semi-Natural Habitats and Field Margins in a Typical Agroecosystem of the Argentinean Pampas as a Reservoir of Carabid Beetles

MARIANA E. MARASAS¹, SANTIAGO J. SARANDÓN^{1,2},
 and ARMANDO CICCHINO³

5

¹Dra. En Ciencias Naturales, Curso de Agroecología, Facultad de Ciencias Agrarias y Forestales, Universidad Nacional de La Plata, Buenos Aires, Argentina

²Ing. Agrónomo. Comisión de Investigaciones Científicas, Pcia. de Buenos Aires, Argentina

³Profesor en Biología. Laboratorio de Artrópodos. Departamento de Biología. Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Buenos Aires, Argentina

10

Changes in edaphic carabidofauna abundance in a wheat crop plot, its field margins, and four semi-natural adjacent habitats were evaluated. A low specific richness of carabids was found in the wheat crop. No species was found exclusively in the wheat plot, but there were species found in the surrounding habitat. The observed responses of different species regarding moisture conditions determined their presence or absence in these semi-natural habitats as well as in dominance structures of each particular ambient. A gradual decrease in the number of captured individuals from the field margin to the center of the wheat plot was observed. Semi-natural habitats and field margins become an important requirement for habitat and shelter of the best represented species of ground beetles, particularly for predatory and omnivorous varieties.

15

20

25

KEYWORDS *specific richness, dominance, boundaries, biodiversity, ecological role*

Address correspondence to Santiago Sarandón Dra. En Ciencias Naturales, Curso de Agroecología, Facultad de Ciencias Agrarias y Forestales, Universidad Nacional de La Plata, CC31, Calle 60 y 119, La Plata (1900), Buenos Aires, Argentina. E-mail: sarandon@ceres.agro.unlp.edu.ar

INTRODUCTION

Increasing concern about the need to preserve biodiversity in agroecosystems has been noted by several authors (Swift et al. 2004, Feehan et al. 2005, Waldhardt et al. 2003, Waldhardt 2003), leading to research into the 35 traits and conditions that allow the maintenance or enhancement of biodiversity. The creation of alternative habitats or the maintenance of undisturbed vegetation assemblages surrounding arable fields has been extensively researched by several authors (Sotherton, 1985; Thomas and Marshall, 1999; French and Elliot, 1999; Dennis et al., 1994; Kromp and 40 Steimberg, 1992; Thomas et al., 1991; Coombes and Sotherton, 1986; Asteraki et al., 2004). Q1

In Argentina, agricultural area with extensive monocultures has increased during recent years. Nevertheless, the grassland Pampean agroecosystems have maintained most of their associated biodiversity. Many areas 45 that have been minimally, if at all, disturbed, and that have been considered predominantly unproductive characterize this region. Moreover, during many years, agronomists have regarded the non-crop areas within agricultural landscapes as hostile and as sources of limitation to production, for example, from weeds, pests, and diseases (Marshall, 2002). Currently, this 50 reductionistic vision is being replaced by a new view that recognizes the ecological role of these undisturbed semi-natural habitats. The challenge is to take advantage of the agroecological benefits gained as a result of the diversity maintained in the spontaneous vegetation composed by native as well as exotic species. As an example, the role played by Carabidae inhabit- 55 ing those habitats in relation to the regulation of pests has been recognized (Landis et al., 2002; Magura, 2002).

From an ecological point of view, agroecosystems must be considered as a complex pattern of cultivated and non-cultivated habitats or patches. This approach can bring valuable information about the behavior of several 60 taxa and may be useful in understanding the role that such biodiversity has in a sustainable agricultural system (Marshall and Moonen, 2002). Landscape structure is a determinant component of the spatial and temporal distribution of some important groups of polyphagous predators, such as ground beetles (Coleoptera, Carabidae) (French and Elliot, 1999; Landis et al., 2002; 65 Swift et al., 2004). This group can be severely affected by agricultural landscape simplification, especially if certain minimal needs of feeding resources, mating sites, refuge,s and adequate habitats for its mobility are limited or scarce (Altieri and Letourneau, 1982; Carmona and Landis, 1999; Portauf et al., 2005). 70

Thomas et al. (1991) found an increase in predator carabid abundance in winter if natural vegetation islands were incorporated into cultivated systems. Different field margins and vegetative groundcover also provide habitats for overwintering of some predatory carabid species that can show a permanent

and cyclic mobility from field margins to cultivated field (Sotherton 1985; 75
Dennis et al., 1994; Thomas and Marshall, 1999; Asteraki et al., 2004).

Q2

In Argentina, important edaphic groups have been identified specifically for their predaceous habits (Marasas et al., 2001; Cicchino et al., 2003) because they play an important role as natural enemies of potential pests of main crops in arable fields. One of the most important families is Carabidae, 80 which is represented by species ranging from medium to large size (2.5 to 25 mm) with cursorial and burrowing behavior (Marasas et al., 2001; Cicchino et al., 2003).

Wheat (*T. aestivum* L.) is among the most important crops in Argentina, and is cultivated in extensive areas (6 millions ha) of landscape homogene- 85 ity that could negatively impact the availability of adequate conditions for survivorship of these ground beetles. Cereal or oilseed monocultures are surrounded by permanent or semi-permanent habitat that increases agricultural landscape heterogeneity. It is thought that adjacent low-disturbed habitats, which are typically grassland of the Pampean agricultural land- 90 scape (Baldi et al., 2006), can play an important role as reservoirs for the best represented predatory carabid species (Cicchino et al., 2003). Nevertheless, these species' abundance can decrease from field margins to the center of the arable fields (Kromp and Steinberger, 1992; Altieri, 1992; French et al., 2001). In order to advance this knowledge, four typical habitats of the 95 Pampean region (larger plains with little drainage) were studied.

The aim of this article is to evaluate species richness, ecological role, dominance structure, and change in the edaphic carabidofauna abundance in a wheat crop plot and its field margins as well as the adjacent semi-natural habitats present in most of the Pampean agroecosystems. 100

MATERIALS AND METHODS

Study Area

Research for this study was carried out at the Experiment Station of the Faculty of Agricultural Sciences, La Plata, Argentina (35° South Latitude). The climate is temperate with mean annual temperature varying between 22 °C 105 for the hottest month (January) and 8 °C for the coldest one (July). Mean annual rainfall varied from 800 to 1000 mm, with no dry season. The field has a typical argiudol soil with some internal drainage deficiencies and the following values at sowing time (0 to 20 cm deep): organic matter 4.2 %, a pH of 6.1, and a C/N ratio of 11.5. 110

Sampling Procedures

Carabid beetles were sampled using pitfall traps. Each trap consisted of plastic pots 150 mm deep, with a diameter of 100 mm, filled with 1/3

volume/volume with a mix of 4% formaldehyde in water, and some drops of detergent as a tensioactive agent. A non-transparent ceramic cover was placed 10 cm above each trap to prevent flooding from rainwater and evaporation of the inside solution. Total number of pitfall traps in each environment was selected according to Obrtel's (1971) criteria. The number of total individuals captured in pitfall traps was evaluated as a measure of the activity-density of the surface-living invertebrates (Thiele, 1977; Baars, 1979).

Two situations were evaluated. The first was conducted during the crop cycle 1996–1997, and the last during 1999. Their main characteristics and conditions were as follows.

Situation 1 was carried out in a patch of 7000 m² cropped with bread wheat (*Triticum aestivum* L.) sown at a density of 120 kg/ha under conventional tillage with mouldboard plough, and fertilized with 100 kg/ha of superphosphate (00-46-00) at sowing. After crop emergence, weeds were controlled with applications of 2-4D plus Picloran at 400 cm³/ha + 100 cm³/ha, respectively. In the crop plot, four strips of 10 m wide per 70 m length each were evaluated. In each strip, three pitfall traps placed 5 m apart from each other were established, resulting in a total of 12 traps. Pitfall traps were collected every 30 to 40 days, totalling 9 samples.

Four semi-natural habitats typical of Argentinean Pampas, located at a distance between 50 and 150 m surrounding cultivated plots, were analyzed:

1. Border of a small lagoon: an artificial and permanent lagoon of 2000 m², which was densely covered with natural vegetation, with a predominance of Poaceae and diverse broad leaf species. The field margins were flooded during times of intensive rainfall.
2. Cortaderia bushes (*Cortaderia selloana* (Sult.) Asch. et Graeb.): this habitat, typical of this region, is characterized by dense shrubs, each of which measure 4 to 10 m². These plant structures provide many microhabitats with relatively stable temperature and humidity conditions.
3. Reed bed: composed of Castilla cane (*Arundo donax* L.) covering a 500 m² area. This stand showed a great density of individuals as well as a thick layer of litter made up of fallen leaves, creating a very homogeneous habitat with stable conditions.
4. Small stand of *Myoporum laetum* Forst. F. of 200 m² composed of sparse low-height trees, the soil surface being covered by a thick layer of litter of dried fallen leaves.

In each of the four semi-natural habitats, 3 plots 5 m apart from each other were established. In each one, 3 pitfall traps were placed, resulting in a total of 9 traps per semi-natural habitat.

Situation 2 was carried out in a crop plot of 10,000 m². Half of the plot was cultivated with bread wheat (*Triticum aestivum* L.) under conventional

tillage with mouldboard plough (treatment A). The other half was maintained without crop during the whole sampling period (treatment B) by means of frequent tillage. The crop plot was surrounded by margins of a ridge 0.5 m high and 2.00 m wide, covered mainly by broad leaf annual species (*Brassica sp.*, *Capsella bursa-pastoris* (L.) Medikus, *Taraxacum officinale* Web. and *Trifolium sp.*) and grasses such as *Bromus unioloides* H.B.K. and *Avena sp.* Three parallel transects were established across the plot at center, 10 m apart from each other from one field margin to another. In each transect, pitfall traps were placed 10 m apart, totalling 17 traps per transect. Pitfall traps were placed on September 18 and were collected 4 times during each 30-day period during spring, which had been determined to be the period of highest activity of the studied species (Marasas et al., 1997) and is in accordance with the results of Biaggini et al. (2007).

Material Identification and Data Processing

In both experiments, all collected individuals were identified at the species level. Carabidae number was recorded and a dominance structure in crop plots and in semi-natural habitats was constructed, based on the proportional distribution of the individuals per species over the total number of individuals sampled. Categories cited by Agosti and Sciaky (1998) were used, as follows: Eudominant: > 10%; Dominant: between 5% and 10%; Subdominant: between 2% and 5%; Recedent: between 1% and 2%; and Subrecedent: < 1%.

Species characterization according to their habitat preferences and for their humidity affinities was done according to Cicchino et al. (2003) and Cicchino and Farina (2005), as follows: Hygrophilic: only tolerate very humid environments, near water bodies Mesophilic: tolerate environments with important humidity variations near or far from water bodies; Xerophilic: live in very open environments and with very low soil and environmental humidity levels.

Individuals belonging to the Carabidae, which is the most abundant and representative family in the Coleoptera order (Marasas et al., 1997; Biaggini et al., 2007), were identified at species level consulting the current bibliography as well as the collections of the Museum of Natural Sciences La Plata and the Museum of Natural Sciences Bernardino Rivadavia of Argentina.

Data were analysed by an unbalanced analysis of variance (ANOVA) considering the mean value of the five traps of each plot. The values were previously transformed by log function. To facilitate interpretation of the data, figures were presented untransformed. A regression analysis between some variables was done. Differences of means were determined by LSD test at $p < 0.05$.

RESULTS

Situation 1: During the entire sampling period (June 1996 to January 1997), a total of 1567 individuals of Carabidae were captured and grouped according to their humidity affinity, habitat preference and relative dominance (Table 1). Ten species were exclusively found in semi-natural habitats: *Selenophorus alternans* (Dejean, 1829), *Notaphus (N) laticollis* (Brullé, 1838), *Notiobia (Anisotarsus) cupripennis* (Germar, 1824), *Brachinus (Neobrachinus) pallipes* (Dejean, 1826), *Incagonum lineatopunctatum* (Dejean, 1831), *Metius circumfusus* (Germar, 1824), *Stenocrepis (Stenocrepis) laevigata* (Dejean, 1831), *Polpochila (P) pueli* (Negrè, 1963), *Paratachys bonariensis* (Steinheil, 1869), and *Apenes cfr. Erythrodera* (Chaudoir, 1875). None of these were found in the wheat plot during the entire sampling period.

For all of the species, a similar proportion of groups of humidity affinity were found between crop plot and *Cortaderia* habitat (60% vs. 65% of mesophilic, 35% vs. 35% hygrophilic, and 5% vs. 0% xerophilic). In the border of the lagoon, a dominance of hygrophilic over the mesophilic species (58.4% vs. 41.6%) was observed. In the crop plot, all of the dominant species were of predatory behavior.

In the reed bed and stand of *Myoporum* habitats, a lower number of carabid individuals was observed (Figure 1) in comparison to the borders of small lagoon and *Cortaderia* bush habitat ($F = 11.87$; $p = 0.000$). Thus, only these last two habitats were compared with the cultivated one. Within these three habitats, only those species whose dominance was equal or higher than recedent were analyzed.

Species richness and dominance structure of carabidae showed differences among habitats. All of the eight species considered relevant according to their dominance in the wheat crop plot (Table 1) were present in lagoon borders, and of those, five were also found in the *Cortaderia* bush. No exclusive species were found in wheat plots. The most represented species were *Paranortes cordicollis* (Dejean, 1828), *Scarites (Scarites) anthracinus* (Dejean, 1831), *Aspidoglossa intermedia* (Dejean, 1831), and *Pachymorphus striatulus* (Fabricius, 1792). All of them were ubiquitous eurytopic and synanthropic species (Cicchino, 2003; Cicchino et al., 2005). Other species, such as *Metius circumfusus* (Germar, 1824), were only found in the lagoon borders and *Cortaderia* bush and with very different dominance, and *Bradycellus sp* was only found in this latter habitat. These last two hygrophilic species have a marked tendency to omnivorism (Cicchino and Farina, 2005).

Carabid abundance in the wheat crop plot, lagoon borders and *Cortaderia* bush, showed seasonal fluctuations (Figure 2). At the beginning of the sampling period, during autumn and winter months, just after wheat was sown (samplings 1, 2, 3, 4, and 5), the highest abundance was found in

TABLE 1 List of Species Captured in Wheat Crop Plot, Lagoon Borders, *Cortaderia* Bush, Reedbed and Small Stands of *Myoporum laetum*; Habitat Preference, Affinities for Humidity (Cicchino and Farina, 2005) and Dominance Structure of the Argentinean pampa: (SubR: subprecedent; R: recedent; SubD: subdominant; D: dominant; EuD: eudominant)

Species	Habitat preference	Affinities humidity	Wheat crop		Lagoon borders		Cortaderia bush		reedbed		<i>Myoporum laetum</i>	
			(n%)	Dominance	(n%)	Dominance	(n%)	Dominance	(n%)	Dominance	(n%)	Dominance
<i>Pachymorphus striatulus</i> (Fabricius, 1792)	Eurytopic	Mesophilic	7	D	3.5	SubD	8.1	D	14.7	EuD	0.63	SubR
<i>Pachymorphus</i> sp.	Stenotopic	Mesophilic	0.39	SubR	0.13	SubR	1	R				
<i>Paranortes cordicollis</i> (Dejean, 1828)	Eurytopic	Mesophilic	23.9	EuD	21.5	EuD	47.3	EuD	13.6	EuD	2.5	SubD
<i>Selenophorus anceps</i> (Dejean, 1831)	Eurytopic	Mesophilic	0.39	SubR	0.34	SubR	8.1	D	0.98	R		
<i>Selenophorus alternans</i> (Dejean, 1829)	Eurytopic	Mesophilic					2.3	SubD				
<i>Selenophorus punctulatus</i> (Dejean, 1829)	Stenotopic	Xerophilic	0.39	SubR								
<i>Notaphus (N.) laticollis</i> (Brullé, 1838)	Stenotopic	Hygrophilic			0.34	SubR						
<i>Aspidoglossa intermedia</i> (Dejean, 1831)	Eurytopic	Hygrophilic	33.7	EuD	9.2	D	0.3	SubR	1.96	R	6.3	D
<i>Scarites (S.) melanarius</i> (Dejean, 1831)	Eurytopic	Mesophilic	0.39	SubR	1.3	R	3.4	SubD				
<i>Scarites (S.) anthracinus</i> (Dejean, 1831)	Eurytopic	Mesophilic	19.6	EuD	11.9	EuD	0.67	SubR				
<i>Clivina (Paraclivina) breviscula</i> (Putzeys, 1866)	Eurytopic	Hygrophilic	5.49	D	0.34	SubR						
<i>Clivina (Paraclivina) media</i> (Putzeys, 1866)	Eurytopic	Mesophilic	0.78	SubR	0.34	SubR						
<i>Bradycellus</i> sp.	Eurytopic	Hygrophilic	0.39	SubR	2.2	SubD	0.3	SubR			0.63	SubR

(Continued)

TABLE 2 (Continued)

Species	Habitat preference	Affinities humidity	Wheat crop		Lagoon borders		Cortaderia bush		reedbed		<i>Myoporum laetum</i>	
			(n%)	Dominance	(n%)	Dominance	(n%)	Dominance	(n%)	Dominance	(n%)	Dominance
Loxandrus simplex (Dejean, 1831)	Eurytopic	Hygrophilic	0.39	SubR	0.13	SubR	1	R	0.98	SubR	3.2	SubD
Loxandrus confusus (Dejean, 1831)	Eurytopic	Hygrophilic	1.96	R	2.53	SubD	1	R	1	R	3.2	SubD
Notiobia (Anisotarsus) cupripennis (Germar, 1824)	Eurytopic	Mesophilic					1.7	R				
Galerita collaris (Dejean, 1826)	Eurytopic	Mesophilic	0.78	SubR	0.34	SubR	0.67	SubR	3.9	SubD	0.63	SubR
Calosoma retusum (Fabricius, 1775)	Eurytopic	Mesophilic	0.39	SubR					0.98	R		
Argutoridius bonariensis (Dejean, 1831)	Eurytopic	Mesophilic	1.56	R	4	SubD	8.8	D	24.5	EuD	85	EuD
Argutoridius chilensis ardens (Dejean, 1828)	Eurytopic	Hygrophilic	1.17	R	3.4	SubD	12.2	EuD	1	R	1.3	R
Brachinus (Neobrachinus) pallipes (Dejean, 1826)	Stenotopic	Hygrophilic			1.7	R						
Incagonum lineatopunctatum (Dejean, 1831)	Eurytopic	Mesophilic					1	R	1.96	R	1.3	R
Metius circumfusus (Germar, 1824)	Euritopics	Hygrophilic			33.5	EuD	2	SubD	4.9	SubD		

Stenocrepis (Stenocrepis) laevigata (Dejean, 1831)	Stenotopic	Hygrophilic	0.92	R	2.9	SubD
Pericompsus (Eidocompsus) metallicus (Bates, 1871)	Stenotopic	Hygrophilic	0.39	SubR		
Pericompsus (Pericompsus) callicalymma (Erwin, 1974)	Stenotopic	Mesophilic	0.39	SubR	0.3	SubR
Polpochila (P) pueli (Negre, 1963)		Hygrophilic	0.13	SubR	0.3	SubR
Paratachys bonariensis (Steinheil, 1869)	Stenotopic	Hygrophilic	0.4	SubR		
Apenes cfr. Erythrodera (Chaudoir, 1875)		Mesophilic				0.63 SubR
Pelmatellus sp.		Mesophilic				0.63 SubR
Total number (1567)			757		102	157




FIGURE 1 Number of individual ground beetles captured per trap in crop plots, lagoon borders, *Cortaderia* bush, reed bed and *Myoporum laetum* stands. Letters above bars indicate significant differences according to LSD Test at $p = 0.05$.




FIGURE 2 Seasonal changes in the abundance of the Carabidae family in different habitats: wheat crop plots, Lagoon borders and *Cortaderia* bush during all sampling period. Vertical lines represented mean standard error.

lagoon borders and *Cortaderia* bush habitat. In spring (sampling 6, 7, 8, 240 and 9), a gradual increase of carabid individuals in the crop field was observed, particularly in comparison to the *Cortaderia* bush habitat.

Situation 2: A total number of 306 individuals were captured over the whole sampling period (September 18 to January 18). No significant differences were found between abundance and the species richness of 245 individuals captured in both field margins.




FIGURE 3 Relationship between the distance from field margin and number of individual Carabidae. Vertical lines represented mean standard error.

Within Carabidae species, the most representative were *Scarites* (*S*) *anthracinus*, *Aspidoglossa intermedia*, *Parypathes* (*P*) *cordicollis* and *Pachymorphus striatulus*. In the field-margins, the fosor and semifosor species (*Scarites anthracinus* and *Aspidoglossa intermedia*) account for 80% of 250 the captured individuals. Total number of individuals captured in A and B treatments were similar.

A marked tendency toward a gradual decrease in total number of individuals from field margin to the center of the plot was observed in A treatment (wheat crop). A negative and significant relationship ($R^2 = 0.7279$ 255 $**p < 0.01$) between the number of individuals and the distance from the field margin was observed. The number of individuals captured 50 m away from the field margin was significantly lower than those found in the margin (Figure 3). In contrast, no significant correlation was found between the distance from the margin and the number of individuals captured in the center 260 of the plot in treatment B.

DISCUSSION

The ecological role of field margins is now being reconsidered, mainly in relation to agrobiodiversity conservation (Marshall and Moonen, 2002). The old vision that field margins constituted a barrier to the expansion of 265 cropped areas is now changing, and the role of field margins as refuges for important predatory and polyphagous carabids is now being recognized (Marshall, 2002; Vanbergen et al., 2005). The conservation of non-disturbed field margins in crop fields would preserve these functional groups and could promote the sustainability of agroecosystems. 270

Results presented here suggest the importance of semi-natural reservoirs of vegetation for extensive cropping systems, such as the monoculture of cereal crops in Argentinean pampas. These habitats can act as refuges for predatory carabids, as well as for omnivorous soil arthropods, and can, in this way, mitigate the effects of monocultures (Sotherton, 1985; Dennis et al., 1994; Thomas and Marshall, 1999; Asteraki et al., 2004). Presence of such functional groups will help to keep phytophagous populations under control and to regulate insect pest outbreaks (Portauf et al., 2005; Swift et al., 2004; Fournier and Loreau, 2002; Krooss and Schaefer, 1998; Edwards et al., 1979).

These semi-natural habitats guarantee the conservation of stable arthropod communities (Woodcock et al., 2005). These are associated with plant species richness and landscape architecture at a local scale and with the conservation of habitats with different degrees of humidity (Gudleifsson, 2005; Cicchino et al., 2005).

It was observed that species had different habitat preferences. The higher number of species found in semi-natural habitats (lagoon borders and *Cortaderia* bush) in relation to wheat crop plots suggests the existence of several microhabitats associated with structural and functional diversity of these habitats (Swift et al., 2004). Lagoon margins are characterized by a high floristic richness and high plant density. Woodcock et al. (2005) and Asteraki et al. (2004) pointed out that non-disturbed vegetative diversity provides ecological niches that harbor a significant abundance and variety of predaceous species. Nevertheless, in the *Cortaderia* bush, the existence of microhabitats was determined, not by plant-specific diversity, but by the shrub structure itself, which has multiple vertical layers that create conditions of shadow and humidity favoring the permanence of carabid communities.

Lower abundance and a lower specific carabid richness were observed in arable field as compared to semi-natural habitats. This could be due to the existence of a reduced quantity of available microhabitats because of the system's homogeneity, as well as changing conditions due to the high disturbance associated with crop management techniques (Gudleifsson, 2005).

The observed responses of different species regarding humidity conditions (hygrophilic, mesophilic, and xerophilic) determined their presence or absence in these habitats and their dominance levels (Table 1). This was reflected in the high proportion of hygrophilic species found both in *Cortaderia* bush and lagoon borders habitats, which suggests that they provide necessary conditions for its permanence in the agroecosystem.

The fact that a non-exclusive carabid species was found in wheat crop plots, and a greater abundance of carabids was found in semi-natural habitats, suggests that the latter can act as a refuge and provide sites for reproduction and hibernation (Pfiffner and Luka, 2000). Landscape simplification

because of crop monoculture would lead to a loss of these habitats and 315
their associated species, increasing the risk of their disappearance in these
agroecosystems. Dominant species in the crop field were also the domi-
nant species in lagoon borders and *Cortaderia* bush. These semi-natural
habitats, little disturbed, could act as reservoirs of a great percentage of
predator carabids and could exert a strong influence on crops. These hab- 320
itats would act as permeable boundaries from and to the field crop and
would therefore be an efficient mode of colonizing predator carabid
individuals. (Lopez Barrera, 2004). The seasonal variation in carabid-
ofauna abundance found both in crop and in semi-natural habitats sug-
gests that there could be a migration of carabids between the crop and 325
those semi-natural habitats.

Field crop margins also provide adequate habitats to guarantee per-
manent mobility of the carabidofauna between these habitats (Sotherton,
1985; Dennis et al., 1994; Thomas and Marshall, 1999; Asteraki et al.,
2004). The observed differences in the behavior of burrowing species as 330
compared with those with cursorial habits confirm the existence of a
close relationship between habitat characteristics and carabid behavior.
The number of burrowing species (*Scarites* genus) found in field mar-
gins was higher than that found in crop plots. Marasas et al. (1997)
found that such species prefer these low or non-disturbed habitats (like 335
field margins and crops under no till management) to build their colo-
nies. The behavior of the cursorial species was more variable and erratic
because their presence is more conditioned by their search for food
(Fournier and Loreau, 2002).

The localization of field margins as compared to field crops is an 340
important fact that must be taken into account in the design of sustainable
agroecosystems. Our study showed a decrease in carabidae abundance
from field margins to the center of the plot cultivated with wheat (Treatment
A), and is similar to that found in cultivated plots of temperate regions
(Kromp and Steinbeger, 1992; Altieri, 1992; French et al., 2001). This can 345
probable be related to the higher uniformity of the crop landscape that
causes a low mobility of those species of Carabidae of big size as these
seems to be more vulnerable to predation in open spaces or fields (Brose,
2003). Nevertheless, this decrease in carabid number toward the center of
the plot did not occur in the other half of the plot (treatment B) which was 350
maintained without crop throughout the entire sampling period, which sug-
gests that under different conditions, other variables could be important in
an analysis of the influence of field margins. These results confirm the
importance of patterns of land-use in beetle community structure (Vanbergen
et al., 2005). Those landscape characteristics that favor appropriate habitats 355
for shelter, moving, feeding, as well as other landscape characteristics that
guarantee the permanence and activity of carabid species, must be
enhanced or recreated.

ACKNOWLEDGMENTS

We thank Guillermo Cap for comments on a previous version of the manuscript. This work was supported by La Plata University and formed part of the Scientific Research Program.

REFERENCES

- Agosti M. and R. Sciaky. 1998. Carabidocenosi dei vigneti: rapporti con le zone limítrofe ed evoluzione nel tempo. *Natura Bresciana, Ann. Mus. Civ. Sc. Nat., Brescia* 31: 69–86. 365
- Altieri M.A. and D.K. Letourneau. 1982. Vegetation managment and biological control in agroecosystems. *Crop Protection* 1: 405–430.
- Altieri M.A. 1992. *Biodiversidad, agroecología y manejo de plagas*. CETAL (Centro de Estudios de Tecnologías Apropriadas para América Latina y el Caribe). Chile. 162 pp. 370
- Asteraky E.J., B.J. Hart, T.C. Ings, and W.J. Manley. 2004. Factors influencing the plant and invertebrate diversity of arable field margins. *Agriculture Ecosystems and Environment* 102: 219–231.
- Baars M.A. 1979. Catches in pitfall traps in relation to mean densities of carabid beetles. *Oecología* 41: 25–46. 375
- Baldi G.J., J.P. Guershman, and J.M. Paruelo. 2006. Characterizing fragmentation in temperate South America grassland. *Agriculture, Ecosystems and Environment* 116:197–208.
- Biaggini M.R., L. Consorti, M. Dapporto, E. Dellacasa, D. Paggetti, and C. Corti. 2007. The taxonomic level order as a possible tool for rapid assessment of Arthropod diversity in agricultural landscapes, *Agriculture Ecosystems Environment* 122 (2):183–191. 380
- Brose U. 2003. Bottom-up control of carabid beetle communities in early successional wetlands: mediated by vegetation structure or plant diversity?. *Oecologia*, 135: 407–413. 385
- Carmona, D. and D. Landis. 1999. Influence of refuge habitats and cover crops on seasonal activity-density of ground beetles (Coleoptera: Carabidae) in field crops. *Environmental Entomology*. 28: 1145–1153.
- Cicchino A., M.E. Marasas, and M.F. Paleólogos. 2003. Características e importancia de la carabidofauna edáfica de un cultivo experimental de trigo y sus bordes con vegetación espontánea en el partido de La Plata, Pcia. de Buenos Aires. *Revista de Ciencias y Tecnología de la UNSE*. 8: 41–55. 390
- Cicchino A.C. 2003. La carabidofauna edáfica de los espacios verdes del éjido urbano y suburbano marplatense. Su importancia como herramienta de manejo de estos espacios. *Revista de Ciencia y Tecnología, Facultad de Agronomía, UNSdE*. 8: 145–164. 395
- Cicchino A.C., M.E. Marasas, and M.F. Paleólogos. 2005. Fenología y densidad-actividad de cinco especies de Carabidae (Coleoptera) edáficas en un cultivo experimental de trigo y su entorno en el Partido de La Plata, Provincia de Buenos Aires. In:

- V Reunión Científico Técnica de Biología del Suelo y V Encuentro sobre Fijación Biológica de Nitrógeno, Área Temática I, Comunidades Terrestres*, I: 1–14. 400
- Cicchino A.C. and J.L. Farina. 2005. Carabidofauna de los suelos linderos con la Laguna Litoral de la Reserva Natural del Puerto de Mar del Plata, Provincia de Buenos Aires, Argentina. In: *V Reunión Científico Técnica de Biología del Suelo y V Encuentro sobre Fijación Biológica de Nitrógeno, Área Temática I, Comunidades Terrestres*, III: 1–15. 405 **Q6**
- Coombes D.S. and N.W. Sotherton. 1986. The dispersal and distribution of polyphagous predatory Coleoptera in cereals. *Annals of Applied Biology* 108: 461–474.
- Dennis P., M.B. Thomas and N.W. Sotherton. 1994. Structural features of field boundaries which influence the overwintering densities of beneficial arthropod. *Journal of Applied Ecology*. 31: 361–370. 410
- Edwards C.A., K.D. Sunderland and K.S. George. 1979. Studies on polyphagous predators of cereal aphids. *Journal of Applied Ecology*, 16: 811–823.
- Feehan J., D.A. Gillmor, and N. Culleton. 2005. Effects of an agri-environment scheme on farmland biodiversity in Ireland. *Agriculture, Ecosystems and Environment* 107 (2–3): 275–286. 415
- Fournier, E. and M. Loreau. 2002. Foraging activity of the carabid beetle *Pterodictus melanarius* III. In field margin habitats. *Agriculture, Ecosystems and Environment*. 89: 253–259.
- French B.W. and N.C. Elliot. 1999. Temporal and spatial distribution of ground beetle (Coleoptera: Carabidae) assemblages in grasslands and adjacent wheat fields. *Pedobiologia* 43: 73–84. 420
- French B.W., N.C. Elliot, R.C. Berberet, and J.D. Burd. 2001. Effects of riparian and grassland habitats on ground beetle (Coleoptera: Carabidae) assemblages in adjacent wheat fields. *Environmental Entomology* 30 (2): 225–234. 425
- Gudleifsson B.E. 2005. Beetle species (Coleoptera) in hayfields and pastures in northern Iceland. *Agriculture Ecosystems and Environment* 109: 181–186.
- Kromp B. and K.H. Steinberger. 1992. Grassy field margins and arthropod diversity: a case study on ground beetles and spiders in eastern Austria (Coleoptera: Carabidae; Arachnidae: Aranei, Opiliones). Biotic diversity in agroecosystems. *Agriculture, Ecosystems and Environment* 40: 1–4. 430
- Krooss S. and M. Schaefer. 1998. The effect of different farming systems on epigeic arthropods: a five-year study on the rove beetle fauna (Coleoptera: Staphylinidae) of winter wheat. *Agriculture, Ecosystems and Environment* 69: 121–133.
- Landis D.A., S.D. Wratten and G.M. Gurr. 2002. Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annual Review of Entomology* 45:175–201. 435
- Lopez-Barrera F. 2004. Estructura y función en bordes de bosques. Ecosistemas. Revista Científica y Técnica de ecología y medio Ambiente. Asociación española de ecología terrestre. Año XIII, 1. Enero-abril. (URL:<http://www.aeet.org/ecosistemas/041/revision1.htm>). 440
- Magura T. 2002. Carabids and forest edge: spatial pattern and edge effect. *Forest Ecology and Management* 257:23–37.
- Marasas M.E., S.J. Sarandón, and A.C. Cicchino. 1997. Efecto de la labranza convencional y siembra directa sobre la coleopterofauna edáfica en un cultivo de trigo, en la Pcia. de Bs.As. *Ciencia del Suelo*. 15(2): 59–63. 445

- Marasas M.E., S.J. Sarandón, and A.C. Cicchino. 2001. Changes in soil arthropod functional group in a wheat crop under conventional and no-tillage systems in Argentina. *Applied Soil Ecology* 18: 61–68.
- Marshall E.J.P. 2002. Introducing field margin ecology in Europe. *Agriculture, Ecosystems and Environment* 89: 1–4.
- Marshall E.J.P. and A.C. Moonen. 2002. Field margins in northern Europe: their functions and interactions with agriculture. *Agriculture, Ecosystems and Environment* 89: 5–21.
- Obrtel R. 1971. Number of pitfall traps in relation to the structure of the catch of soil surface Coleoptera. *Acta entomologica Bohemoslovaca* 68: 300–309.
- Pfiffner L. and H. Luka. 2000. Overwintering of arthropods in soils of arable fields and adjacent semi-natural habitats. *Agriculture, Ecosystems and Environment* 78: 215–222.
- Portauf T., I. J. Roschewitz, C. DauberThies, T. Tscharnitke, and V. Wolters. 2005. Landscape context of organic and conventional farms: influences on carabid beetle diversity. *Agriculture, Ecosystems and Environment* 108: 165–174.
- Sotherton N.W. 1985. The distributions and abundance of predatory coleoptera overwintering in field boundaries. *Annals of Applied Biology* 106: 17–21.
- Swift M.J., M.N. Izac, and M. van Noordwijk. 2004. Biodiversity and ecosystem services in agricultural landscapes are we asking the right questions?. *Agriculture Ecosystems Environment* 104: 113–134.
- Thiele H.U. 1977. Carabid Beetles in their environments. Springer-Verlag. Berlin. 369 pp.
- Thomas M.B., S.D. Wratten, and N.W. Sotherton. 1991. Creation of island habitats in farmland to manipulate populations of beneficial arthropods: predators densities and emigration. *Journal of Applied Ecology*. 28: 906–917.
- Thomas M.B. and E.J.P. Marshall. 1999. Arthropod abundance and diversity in differently vegetable margins of arable fields. *Agriculture, Ecosystems and Environment*. 72: 131–144.
- Vanbergen, A.J., B.A. Woodcock, A.D. Watt, and Niemela J. 2005. Effect of land-use heterogeneity on carabid communities at the landscape scale. *Ecography* 28: 3–16.
- Waldhardt R. 2003. Biodiversity and landscape- summary, conclusions and perspectives. *Agriculture, Ecosystems and Environment* 98:305–309.
- Waldhardt R., D. Simmering, and H. Albreecht. 2003. Floristic diversity at the habitat scale in agricultural landscapes of Central Europe. Summary, conclusions and perspectives. *Agriculture, Ecosystems and Environment* 98: 79–85.
- Woodcock B.A., D.B. Westbury, S.G. Potts, S.J. Harris, and V.K. Brown. 2005. Establishing field margins to promote beetle conservation in arable farms. *Agriculture, Ecosystems and Environment* 107 (2–3): 255–266.